

# Schizophrenia (Children and Adolescents) - Indicators and Standards

Indicator Domain	Indicator	Type	Standard	Time Reference
<b>Diagnostic Proce</b> s	Proportion of incident (recently diagnosed) patients with schizophrenia, examined for  a. Psychopathology b. Cognitive function c. Need for social support d. Duration of untreated psychosis	Process	a.1. $\geq 98\%$ (psychopathology)  a.2. $\geq 80\%$ (assessed using a diagnostic instrument) (SCAN, OPCRIT, SCID, KIDDI-SADS)  b. $\geq 50\%$  c. $\geq 80\%$  d. $\leq 50\%$	a1: During hospitalisation  a2: During hospitalisation  b: During hospitalisation c: During hospitalisation d: Not exceeding six months in the initiation of treatment after diagnosis
<b>Contact</b>	Proportion of patients with schizophrenia in contact with the treatment system	Process	$\geq 85\%$	No time reference, all patients
<b>Contact person</b>	Proportion of patients with schizophrenia in contact with a specific member of the therapist team or for whom a psychotherapeutic course of treatment is implemented	Process	$\geq 95\%$	No time reference, all patients
<b>Medication</b>	Proportion of patients with schizophrenia on anti-psychotic medication	Process	a. $\geq 90\%$ (Antipsychotics) b. $\geq 80\%$ (Second Generation Antipsychotics) (SGA)	No time reference, all patients
<b>Side effects</b>	Proportion of patients with schizophrenia examined for side effects of the medicine	Process	$\geq 98\%$	No time reference, all patients
<b>Family Intervention</b>	Proportion of patients with schizophrenia where relatives accept the offer of contact with the treatment system	Process	$\geq 98\%$	No time reference, all relatives

<b>Psycho education (education about disease)</b>	<p>a. Proportion of children and adolescents accepting the offer of psycho-education individually</p> <p>b. Proportion of children and adolescents accepting the offer of psycho-education in manualized courses</p> <p>c. Proportion of children, adolescents and relatives who are offered psycho-educational family intervention</p>	Process	<p>a. <math>\geq 90\%</math> (received psycho education, individual sequences)</p> <p>b1. <math>\geq 40\%</math> (patients with incident schizophrenia, manualized courses)</p> <p>b2. <math>\geq 20\%</math> (patients with prevalent schizophrenia, manualized courses)</p> <p>c1. <math>\geq 70\%</math> (relatives to children and adolescents with incident schizophrenia)</p> <p>c2. <math>\geq 50\%</math> (relatives to children and adolescents with prevalent schizophrenia)</p>	<p>a. b. and c: No time reference, all patients</p>
<b>Planned outpatient treatment by discharge</b>	<p>Proportion of patients with schizophrenia participating in planned support measures after discharge (GAPD score <math>\geq 5</math>)</p>	Process	<p>a. <math>\geq 90\%</math> (planned support measures)</p> <p>b. <math>\geq 90\%</math> (receive psychiatric post-discharge treatment)</p> <p>c. <math>\geq 90\%</math> (status for outpatients within 1½ year)</p>	<p>a. and b: No time reference, all patients</p> <p>c: Within 1½ year</p>
<b>Prevention of Suicide</b>	<p>Proportion of patients with schizophrenia assessed for suicide risk at discharge (documented in the patient record and documented in the patient record)</p>	Process	<p><math>\geq 90\%</math></p>	<p>At discharge</p>